

CHEMICAL INVESTIGATION OF ESSENTIAL OIL OF POLYGONATUM VERTICILLATUM (LILIACEAE)

*P.S. Rajpoot,

Associate Professor

Laboratory of Natural Product

Department of Chemistry,

Govt. P.G. College, Chharra, Aligarh (U.P.) 202130 (INDIA)

Email: rajpootphoolsingh@gmail.com

KEY WORDS: IDEX PolygonatumVerticillatum, Liliaceae, Polygonatum
Rhyzom, Essential Oil Composition, Natural Fatty
Acids & Esters

INTRODUCTION :

PolygonatumVerticillatum (Liliaceae) is an erect glabrous natural herb growing at an altitude of 9000-12500 ft. of Kumaon Himalayan Mountains in Uttaranchal, India. The root stock of the plant is sweet, cooling, emollient, diuretic, aphrodisial, galactagoga, aspestising. The plant belongs to Astvag group medicinal Plants rhizomes used is as an ingredient of Chyvanprash and an important Ayurvedic tonic. It is useful in vitiafed conditions of Pitta and Vata, Burning sensation, fever, strangury seminal weakness, female weakness and in problems of reproductive systems. Literature search revealed that glucose,

galactose, four saponosides Lectins, Lysine, Serine, aspartic acid reported from this plant and suggested this plant as a new source for diosgenin production. As part of a search for useful high altitude Himalayan herb this plant was collected and chemical investigated. There is no literature report on essential oil of P. Verticillatum so forth this is the first work essential oil of P. Verticillatum and first time and first time the plant has been found to be rich in natural long chain fatty acids and esters.

RESULTS AND DISCUSSION :

The compounds of essential oil of P. Verticillatum were identified on the basis of their GC/GC-MS analysis and by comparison of their mass spectra with their authentic compounds existing in literature. The volatile constituents identified in the essential oil of P. Verticillatum along with their retention times, percentage areas mass spectra are given in Table-1. The compounds are listed according to their retention time from DB-5 Column. The oil was rich with long chain fatty acids and esters. Lauric acid, Myristic acid palmitic acid, B-Methyl, Metylester and Hexadecanoic acid present in the oil.

Lauric acid or dodecanoic acid – $\text{CH}_3\text{-(CH}_2\text{)}_{10}\text{-COOH}$,

Myristic acid or Tetradecanoic acid – $\text{CH}_3\text{-(CH}_2\text{)}_{12}\text{-COOH}$,

Pentadecanoic acid $\text{CH}_3\text{-(CH}_2\text{)}_{13}\text{-COOH}$,

Pentadecanoic acid 13-methyl-methylester $\text{CH}_3\text{-(CH}_2\text{)}_{15}\text{-COOH}$,

Polmitic acid or n-hexadecanoic acid $\text{CH}_3\text{-(CH}_2\text{)}_{14}\text{-COOH}$

EXPERIMENTAL :

Plant Material : The whole plant *P. Verticillatum* (Liliaceae) was collected the month of October 1998-2003 from Munsyari region of Kumaon Himalaya, Uttaranchal, India at an altitude of 11000-125000 ft. the plant was identified in the Department of Botany, Kumaon University, Nainital as well as at Forest Research Institute, Dehradun.

Extraction of Oil : Rhizome of *P. Verticillatum* (500gm.) was used for essential oil extraction by Clevenger's apparatus the condensate was treated with n-Hexane. The n-Hexane. The hexane layer separated. The organic phase was dried over Na_2SO_4 and the solvent was evaporated, under reduced pressure in a thin film rotary evaporator at 30°C yield of the oil was 3 ml.

GC Analysis : The oil was analysed by GC using flame ionization detector (FID). The temp. program $60\text{-}240^\circ\text{C}$ at $3^\circ\text{C}/\text{min}$. injector temp. 240°C , detector temp. 280°C , total run time 35 min.

GC-MS Analysis : GC-MS was done using fused silica gel capillary column (30x0.25mm) liquid phase DB-5 with helium as a carrier gas in a Hewlett-Packard 5840. GC interfaced with Hewlett-Packard 5985 mass spectrometer. The Column Temperature was programmed at $3^\circ\text{C}/\text{min}$. from $60^\circ\text{-}240^\circ\text{C}$, the analysis by electron impact, 70 eV.

Table - 1 GC-MS of the Oil of Rhizome of *P. Verticillatum*

Sr.No.	Ref.Time	Area%	[M ⁺] Mad spectra	Compounds
1.	19.44	13.0%	200	Lauric Acid
2.	24.96	40.0%	228	Myristic Acid
3.	27.36	25.0%	242	Pentadecyclic Acid
4.	28.82	12.0%	270	13-Methyl-Methyester
5.	30.14	88.0%	256	Polmitic Acid

Table-1 reports the result of the essential oil of P.Verticillatum shade dried rhizom.

Acknowledgement : We thank to RSIC and CDRI Lucknow for GC-MS studies and

Prof. K.S. Khetwal, Deptt. of Chemistry, Nainital University.

REFERENCES –

1. Chopra R.N., Nayar, S.L. and Chopra, I.C. “Glossary of Indian Medicinal Plant” CSIR, New Delhi, 1956.
2. Janeczko Z., Sibiga A, Herba Pol, 1982, 28 (3-4), 11-22.
3. Khetwal, K.S., Verma, D.L., Pathak, R.P., et.al. Indian drugs, 1985, 2(3) 126-128.
4. Koves, Margit, Penksza, Karoly, Turesangi, Gabor, Siller, Iren, Kaszab, Haszlov; Verh Gas Oekol 1996, 25, 147-152
5. Rastogi, R.P. and Mehrotra B.N., “Compendium of Indian Medicinal Plants” Vol.3 1980-1984
6. Shankar, V.Prakas S; Unijal M.R., Nature, Appl., Sci. Bull 1970, 22 (3-4) 139.
7. Albert, Y. Leung. (John Willy and Sons, New York) Encyclopedia of common natural ingredients used in food and cosmetics 1980.
8. Antonjuek, V.O. etal, 1993.
9. Chopra, R.N. Nayr, S.L. and Chopra, I.C. Indogenous drugs of India 1958.
10. Davidson, S.R., Passmore, J.F. Brock and A.S. Truswell (eds) “Human Nutrition and dieties” 6thedi. Churchill Livingstone Edinbarg 1972, 1975.